



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Frontal recession of Gangotri Glacier, Garhwal Himalayas, from 1965 to 2006, measured through high-resolution remote sensing data

Bhambri, R ; Bolch, T ; Chaujar, R K

Abstract: We report in this communication the length fluctuation and frontal area changes at the snout of Gangotri Glacier based on high-resolution satellite data from 1965 to 2006. Glacial outlines were mapped from declassified imageries from Corona (1965, 1968), Hexagon (1980) and Indian satellites IRS PAN (2001) and Cartosat-1 (2006). The results show that Gangotri Glacier exhibited retreat up to 819 ± 14 m and lost 0.41 ± 0.03 sq. km (0.01 sq. km year⁻¹) at its front from 1965 to 2006. The retreat rates are lower than those previously reported using coarse-resolution remote sensing data and the Survey of India topography map. The results of the present study are supported by in-situ field survey conducted by the Geological Survey of India.

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-59630>

Journal Article

Published Version

Originally published at:

Bhambri, R; Bolch, T; Chaujar, R K (2012). Frontal recession of Gangotri Glacier, Garhwal Himalayas, from 1965 to 2006, measured through high-resolution remote sensing data. *Current Science*, 102(3):489-494.

18. Sridhar, A. R., Prasad, A. S. S. R. S., Satyavani, N. and Sain, K., Sub Trappean Mesozoic sediment in the Narmada basin based of travel time and amplitude modelling – a revisit to old seismic data. *Curr. Sci.*, 2009, **97**(10), 1462–1466.
19. Sridhar, A. R., Tewari, H. C., Vijaya Rao, V., Satyavani, N. and Thakur, N. K., Crustal velocity structure of Narmada Son Lineament along the Thuadara–Sendhwa–Sindad profile in the NW part of Central India and its Geodynamic implications. *J. Geol. Soc. India*, 2007, **69**, 1147–1160.
20. Krishna, V. G., Kaila, K. L. and Reddy, P. R., Synthetic seismogram modelling of crustal seismic record sections from the Koyana DSS profiles in the Western India. In *Properties and Processes of Earth's Lower Crust* (eds Mereu, R. F., St Muller and Fountain, D. M.), American Geophysical Union Monograph 51 IUGG, 1989, vol. 6, pp. 143–157.
21. Krishna, V. G., Kaila, K. L. and Reddy, P. R., Low velocity layers in the subcrustal lithosphere beneath the Deccan Traps region of western India. *Phys. Earth Planet. Inter.*, 1991, **67**(3–4), 288–302.
22. Hirn, A., Prodehl, C. and Steinmetz, L., An experimental test of models of the lower lithosphere in Bretagne (France). *Ann. Geophys.*, 1975, **31**, 517–530.
23. Ansorge, J., Die Feinstruktur des obersten Erdmantels unter Europa und dem mittleren Nordamerika, Ph D thesis, Univ. Karlsruhe, Germany, 1975, p. 111.
24. Faber, S., Refraktionsseismische Untersuchung der Lithosphäre unter den britischen Inseln, Ph D thesis, Univ. Karlsruhe, Germany, 1978, p. 32.
25. Sain, K. and Kaila, K. L., Ambiguity in the solution of the velocity inversion problem and a solution by joint inversion of seismic refraction and wide-angle reflection times. *Geophys. J. Internat.*, 1996, **124**, 215–227.
26. Bottings, A. Y., Hirn, A. and Steinmetz, L., Implications de l'existence d'un canal a moindre vitesse sous Moho. *Bull. Soc. Geol. Fr.*, 1973, **7**(5–6), 500–505.

ACKNOWLEDGEMENT. We thank the Director, CSIR-National Geophysical Research Institute, Hyderabad for encouragement and permission to publish this paper.

Received 30 June 2011; accepted 22 December 2011

Frontal recession of Gangotri Glacier, Garhwal Himalayas, from 1965 to 2006, measured through high-resolution remote sensing data

Rakesh Bhambri¹, Tobias Bolch² and Ravinder Kumar Chaujar^{3,*}

¹Center for Glaciology, Wadia Institute of Himalayan Geology, Dehradun 248 001, India

²Geographisches Institut, Universität Zürich, Switzerland

³Wadia Institute of Himalayan Geology, Dehradun 248 001, India

We report in this communication the length fluctuation and frontal area changes at the snout of Gangotri Glacier based on high-resolution satellite data from 1965 to 2006. Glacial outlines were mapped from de-

classified imageries from Corona (1965, 1968), Hexagon (1980) and Indian satellites IRS PAN (2001) and Cartosat-1 (2006). The results show that Gangotri Glacier exhibited retreat up to 819 ± 14 m and lost 0.41 ± 0.03 sq. km (~ 0.01 sq. km year⁻¹) at its front from 1965 to 2006. The retreat rates are lower than those previously reported using coarse-resolution remote sensing data and the Survey of India topography map. The results of the present study are supported by *in-situ* field survey conducted by the Geological Survey of India.

Keywords: Gangotri Glacier, remote sensing, retreat, satellite data.

GANGOTRI Glacier is the largest glacier (length ~ 30 km) in the Garhwal Himalayas. The Bhagirathi River originates from the snout (Gaumukh; ~ 3950 m asl) of Gangotri Glacier, which is the main source stream of Ganga River (Figure 1). Gangotri Glacier originates from the Chaukhamba group of peaks (~ 6853 – 7138 m asl) and flows northwest towards Gaumukh. About 29% of its total area is covered by debris¹. Gangotri Glacier is one of the well-documented and monitored glaciers in the Indian Himalayas as regards to its snout position. Auden² first systematically mapped the snout and geomorphic features of Gangotri Glacier in 1935 using a plane-table survey at a scale of 1 : 4800. Several scientists from GSI have resurveyed Gangotri Glacier and marked the position of the snout on Auden's plane-table map and measured its length in terms of retreat^{3–6}. Length records though are not the most significant parameter for glacier




Figure 1. Gangotri Glacier with its tributary glaciers. Clean and debris-cover glacier outlines of Gangotri Glacier and tributary glaciers derived from Bhambri *et al.*¹.

*For correspondence. (e-mail: rchaujar@gmail.com)

changes, are easy to measure, and are available for many glaciers around the world⁷. These records are useful to reconstruct glacier mass balance^{8,9}; estimate the glacier contribution to sea-level rise^{10,11}; historical equilibrium-line altitudes¹²; response times of glaciers^{13,14}, and global and hemispheric temperature¹⁵. In addition, several remote sensing-based studies have mapped Gangotri Glacier using various multi-temporal and multi-spectral satellite data and measured its linear retreat^{16–21} (Table 1). Among these studies, few have used Survey of India (SOI) topography maps^{17,18,20} or coarse-resolution satellite data¹⁹ (e.g. Landsat MSS with a spatial resolution of 79 m) to acquire older glacier extents (1960s and 1970s) for quantifying variability records of Gangotri Glacier. However, studies have shown examples of inaccuracies in SOI topographical maps^{22–26}. In addition, interpretation of debris-cover, shadow area and seasonal snow on satellite images is known to be one of the major challenges in glacier inventories and glacier change studies^{26–31}.

Declassified imagery (e.g. Corona and Hexagon) or aerial images from the 1960s and 1970s are ideal to map historic extents of glaciers^{30–35}, and can also be used for comparison with glacier outlines derived from old topography maps^{26,30}. These images are accessible from United States Geological Survey website (USGS; <http://edcns17.cr.usgs.gov/NewEarthExplorer/>). The Corona satellite orbit was near-polar and circular. However, the orbit inclination and equator crossing time were different for different missions³⁴. Therefore, there is no systematic coverage of the earth's surface. Generally, the inclination angle varied from 60° to 100° (measured from the equator)³⁴. However, to the best of our knowledge there have been no attempts earlier to measure length records of Gangotri Glacier from historic high-resolution Corona and Hexagon imageries. Thus, the main goals of the present study are: (1) to generate length records and area vacated by Gangotri Glacier at the snout from declassified imageries, and (2) to compare our results with the SOI topography map (1962) and previous remote sensing and field records.

We used two high-resolution Corona KH4A imageries for 24 September 1965 and 27 September 1968, and KH-9 Hexagon imagery for 8 September 1980 with minimal

snow to extract the historic extent of Gangotri Glacier (Table 2). In addition, orthorectified 2006 high-resolution Cartosat-1 and 2001 IRS-1C PAN images and 2006 ASTER DTM, generated for a previous study³¹ were used for the extraction of recent glacier outlines. The Cartosat-1 imagery was used as base image for rectification of older Corona and Hexagon imageries. These older imageries were co-registered based on a two-step approach: (i) a projective transformation was performed based on ground control points (GCPs) and the ASTER DTM using ERDAS Imagine 9.3, followed by (ii) a spline adjustment using ESRI ArcGIS 9.3 (refs 1, 30). For older declassified imageries between 30 and 50, GCPs were acquired from Cartosat-1 imagery for co-registration.

The glacier extents were manually delineated from panchromatic data of Corona, Hexagon, IRS PAN and Cartosat-1 imagery. For the calculation of length changes, stripes with 50 m distance were drawn parallel to the main flow direction of the glacier (Figure 2). Length change was calculated as the average length from the intersection of the stripes with the glacier outlines³⁶. Based on the outlines of the different years, the area vacated near the snout was also calculated. We also calculated length changes in terms of its retreat along the central flow line to be compared with results derived from average length from the intersection of the stripes with the glacier outlines. Geomorphic features such as gully talus and moraines presented by GSI studies^{2,37} in large-scale map were also mapped from 1968 Corona imagery (Figure 3). These moraines were dated using lichenometry³⁷.

Glacier outlines derived from various satellite datasets with different spatial resolutions, obtained at different times with varying snow cover, cloud and shadow conditions have different levels of accuracy. Thus, estimation of the uncertainty is crucial to know about the accuracy and significance of the results. The Spline method

Table 1. Retreat of Gangotri Glacier measured by various authors

Period	Retreat		Reference
	Total (m)	Mean rate per year (m)	
1935–1996	1220	20	6
1962–1999	1250	34	42
1935–1997	2500	40	45
1962–2000	1600	42	17
1985–2001	368	23	21
1962–2000	1510	40	18
1962–2006	1651	38	20





Figure 2. Glacier outlines derived from different satellite imageries and overlaid stripes with 50 m distance. The average retreat rates are derived from the intersection of the glacier outlines with the band of stripes.

Table 2. Details of satellite data used in the present study

Satellite data	Date of acquisition	Spatial resolution (m)	Scene/product ID	Planimetric accuracy
Corona KHA	24 September 1965	4	DS1024-1023DF117	± 5.2
Corona KHA	27 September 1968	4	DS1048-1134DF107	± 3.2
KH-9 Hexagon	08 September 1980	7	DZB1216-500329L007001	± 12.7
IRS PAN	26 October 2001	5.8	IRS1CDPSR1V4D090002600101	± 10.7
Cartosat-1	28 September 2006	2.5	097001100102	± 10

**Figure 3.** Geomorphic features demarcation on Corona imagery according to Auden² and Vohra³⁷ large-scale map.

provides a precise registration with a minor error¹. To assess the positional accuracy, 11 common geomorphic location points were identified on the base of Cartosat-1 and other Corona KH4A, KH-9 Hexagon and IRS-1C PAN satellite imageries²⁶. The horizontal shift between base Cartosat-1 and corresponding Corona (1965, 1968), Hexagon (1980) and IRS PAN (2001) imageries was found to be 5.2, 3.2, 12.7 and 10.7 m respectively (Table 2). The uncertainty was calculated from the following formula for multi-temporal measures of the glacier front position using satellite images³⁸.

$$e = \sqrt{(a1)^2 + (a2)^2} + E_{\text{reg}},$$

where $a1$ is the pixel resolution of imagery 1, $a2$ the pixel resolution of imagery 2 and E_{reg} is the registration error.

Hence, the uncertainty can be estimated in case of the 1965 Corona imagery as follows:

$$e = \sqrt{[(4)^2 + (2.5)^2]} + 5.2 = 10 \text{ m.}$$

The uncertainty was 8 m for Corona (1968), 20 m for Hexagon (1980) and 17 m for IRS PAN imagery

(2001). The uncertainty for glacial area was estimated by multiplication of the uncertainty of length with glacier width.


Our results show that Gangotri Glacier retreated 819 ± 14 m from 1965 to 2006 (Table 3). On an average, Gangotri Glacier retreated at the rate of 5.9 ± 4.2 m/year from 1965 to 1968 and 26.9 ± 1.8 m/year from 1968 to 1980, and it retreated 21.0 ± 1.2 m/year between 1980 and 2001. The recession rate declined during 2001–2006 and it receded at a rate of 7.0 ± 4.0 m/year. Earlier studies on the recession of Gangotri Glacier using topography map and satellite data show higher estimation of recession than our assessments. For instance, one study²⁰ estimated 1651 m or an average 38 m/year retreat rate of Gangotri Glacier based on the 1962 topography map and 2006 ASTER imagery. Similarly, other studies^{17,18} reported about 1600 and 1510 m recession of Gangotri Glacier at its front from 1962 to 2000 respectively. All these estimations are almost twice as high as our results (1965–2006). However various studies^{18,20} show that the glacier outline derived from the SOI topography map (1962) covers an area near the moraine dated AD 1900 by Vohra³⁷ up to the gully talus^{2,37} (Figure 3). This indicates that higher retreat of Gangotri Glacier is probably

Table 3. Total and average recession of Gangotri Glacier length

Period	Total retreat (m)	Rate of retreat (m/year)	Total length change (m) along the central flow line
1965–1968	-17.7 ± 12.8	-5.9 ± 4.2	$+77 \pm 12.8$
1968–1980	-323.2 ± 21.5	-26.9 ± 1.8	-465 ± 21.5
1980–2001	-441.0 ± 26.2	-21.0 ± 1.2	-537 ± 26.2
2001–2006	-37.0 ± 20	-7.4 ± 4	-6 ± 20
Total	-818.9 ± 14	-19.9 ± 0.3	-1085 ± 14

Table 4. Total and average area vacated near Gangotri Glacier snout

Year	Total area vacated at snout (10^3 m^2)	Average area vacated at snout ($10^3 \text{ m}^2/\text{year}$)
1965–1968	22.9 ± 6.0	7.6 ± 2.0
1968–1980	163.9 ± 10.1	13.7 ± 0.8
1980–2001	215.4 ± 12.3	10.2 ± 0.5
2001–2006	16.39 ± 9.4	3.2 ± 1.8
Total (1965–2006)	418.5 ± 37.8	10.2 ± 0.9

**Figure 4.** Glacier outlines derived from different high-resolution satellite data overlaid on Cartosat 1 (2006) imagery.

associated with the overestimated delineation of the glacier outline on SOI map. We also calculated the recession of Gangotri Glacier as shown in the satellite imagery by Kargel *et al.*³⁹. The older terminus positions shown by Kargel *et al.*³⁹ in satellite imagery are based on

lichnometry^{37,40}. A retreat of $\sim 1444 \text{ m}$ from 1935 to 2006 ($\sim 20.3 \text{ m/year}$) was calculated by Kargel *et al.*³⁹ along the Bhagirathi River, whereas a GSI study⁶ suggests that Gangotri Glacier retreated by $\sim 1220 \text{ m}$ between 1935 and 1996, which amounts to an average rate of $\sim 20 \text{ m/year}$.

We found that Gangotri Glacier receded about 764 ± 19 m between 1968 and 2001 (23.2 ± 0.6 m/year), whereas the field-based GSI study⁶ showed that its lost about 720 m length from 1971 to 1996 (~ 28.8 m/year). Previous remote-sensing studies^{17,18,20} calculated the retreat along the central flow line or more or less along the Bhagirathi River. Our results based on the average length from the intersection of the stripes with the glacier outlines reveal that Gangotri Glacier receded 819 ± 14 m from 1965 to 2006, whereas recession along the central flow line would be 1085 ± 14 m. These results show similar tendencies, but the measure based on one point of the glacier only may overestimate the recession and is more susceptible to outliers. The averaging along the front is a more robust method and provides more reliable estimations, especially in the situation of a change in location of the ice caves (Table 3).

The present study shows that Gangotri Glacier lost 0.41 ± 0.03 sq. km (~ 0.01 sq. km year⁻¹) area between 1965 and 2006 from its front (Table 4). These results are supported by *in-situ* field surveys conducted by GSI⁶. The GSI results indicate that Gangotri Glacier reduced its area at its terminus by 0.58 sq. km (~ 0.01 sq. km year⁻¹) between 1935 and 1996 (ref. 6). In addition, Auden² noticed a single ice cave at the left side of Gangotri Glacier snout. Another GSI study³ reported two caves one small and one large, on the terminus of Gangotri Glacier in 1956. Furthermore, one more field-based photographic evidence shows that the ice near and between the two tunnels of Gangotri Glacier was vertically banded as a result of the flow structure imposed on the glacier where it had reunited after its bifurcation immediately upstream by a rock barrier⁴¹. Another study⁴ noticed only a single large cave on the right fringe of the snout during 1967. However, after 1956, the larger and prominent ice cave had been considered for the recession studies^{4,6}. Shadows of ice caves on the 1965 and 1968 Corona satellite images suggest the existence of two caves at the terminus of Gangotri Glacier (Figure 4). This indicates that high-resolution satellite data provide more reliable results in comparison with previous studies^{17,18,20} based on coarser-resolution satellite data such as Landsat MSS and TM, and/or topography maps. We found slight advancement of the middle part at the terminus of Gangotri Glacier during 1968, which is probably concerned with deposition of ice block of the snout after its disintegration. This natural process was reported by an earlier study⁴². Our results suggest that in recent times (2001–2006) Gangotri Glacier has lost few square metres of the area, which has also been reported by prior field-based studies²⁵. Frontal recession of Gangotri Glacier has shown variability in the amount, rate and time of occurrence during the study period. From 2001 to 2006, the recession of Gangotri Glacier has declined compared to the previous observation during the study period. However, it does not imply that Gangotri Glacier recession has ceased as length changes

show only the indirect and delayed response of a glacier to climate change, in contrast to glacier mass balance. The response time of the large debris-covered Gangotri Glacier is likely to be much longer than that of smaller glaciers in the Garhwal region⁴³. Thus the study of mass balance is needed for precise knowledge of glacier health. However, *in situ* measurements over the entire glacier are logistically difficult and hence not feasible owing to its size and characteristics. The geodetic approach using the presented Corona data⁴⁴ could be a more suitable substitute as well as cost-effective method. Moreover, topographic parameters such as elevation range, aspect as well as glacier size, shape, motion, thickness and distribution of debris-covered area, contribution of tributary glaciers to the accumulation and the local topography influence glacier response, which need to be addressed in further studies.

The main challenges with the Corona data are the complex image geometry and the absence of satellite camera specifications. The positional accuracy of the rectified Corona imagery can be evaluated based on common unchanged location points. We have only addressed the length and frontal glacial area changes measured through the snout position on satellite imageries for different years. The uncertainty is negligible for the small area, but is usually higher when addressing a larger area. The Corona data are useful for providing insight into glacier changes since the 1960s, and the above mentioned field studies^{5,6} corroborate our results for Gangotri Glacier. Suitable declassified images are also available for many glaciers of other remote mountain areas where no aerial images are available or accessible.

1. Bhambri, R., Bolch, T., Chaujar, R. K. and Kulshreshta, S. C., Glacier changes in the Garhwal Himalayas, India 1968–2006 based on remote sensing. *J. Glaciol.*, 2011, **57**(203), 543–556.
2. Auden, J. B., Snout of the Gangotri Glacier, Tehri Garhwal. *Rec. Geol. Surv. India*, 1937, **72**, 135–140.
3. Jangpangi, B. S., Report on the survey and glaciological study of the Gangotri glacier, Tehri Garhwal District: Glacier No. 3, Arwa Valley: Satopanth and Bhagirath Kharak Glaciers, Garhwal District, Uttar Pradesh. *Mem. Geol. Surv. India*, 1958, p. 18.
4. Tiwari, A. P., Study of the Gangotri Glacier, Uttarkashi, Central Himalayas, UP. *Rec. Geol. Surv. India*, 1972, **106**, 248–256.
5. Raina, V. K., Is the Gangotri glacier receding at an alarming rate? *J. Geol. Soc. India*, 2004, **64**, 819–821.
6. Srivastava, D., Recession of Gangotri glacier. In Proceedings of Workshop on Gangotri Glacier, Lucknow, 26–28 March 2003 (eds Srivastava, D., Gupta, K. R. and Mukerji, S.), Geological Survey of India, Special Publication Number 80, 2004, pp. 21–32.
7. Zemp, M. et al. (eds), *Global Glacier Changes: Facts and Figures*, UNEP, World Glacier Monitoring Service (WGMS), Zurich, Switzerland, 2008, p. 88.
8. Hoelzle, M., Haeberli, M., Dischl, M. and Pescke, W., Secular glacier mass balances derived from cumulative glacier length changes. *Gl. Plan. Change*, 2003, **36**, 295–306.
9. Lüthi, M. P., Bauder, A. and Funk, M., Volume change reconstruction of Swiss glaciers from length change data. *J. Geophys. Res. F*, **115**, F04022; doi:10.1029/2010JF001695.

10. Oerlemans, J., Dyurgerov, M. and van de Wal, R. S. W., Reconstructing the glacier contribution to sea-level rise back to 1850. *Cryosphere*, 2007, **1**, 59–65.
11. Leclercq, P. W., Oerlemans, J. and Cogley, J. G., Estimating the glacier contribution to sea-level rise for the period 1800–2005. *Surv. Geophys.*, 2011, **32**, 519–535.
12. Klok, E. J. and Oerlemans, J., Deriving historical equilibrium-line altitudes from a glacier length record by linear inverse modelling. *Holocene*, 2003, **13**, 343–351.
13. Oerlemans, J., Estimating response times of Vadret da Morteratsch, Vadret da Palue, Briksdalsbreen and Nigardsbreen from their length records. *J. Glaciol.*, 2007, **53**(182), 257–362.
14. Pelto, M. S. and Hedlund, C., The terminus behavior and response time of North Cascade glaciers. *J. Glaciol.*, 2001, **47**, 497506.
15. Leclercq, P. W. and Oerlemans, J., Global and hemispheric temperature reconstruction from glacier length fluctuations. *Climate Dyn.*, 2011; doi:10.1007/s00382-011-1145-7.
16. Chaujar, R. K., Mazari, R. K. and Gergan, J. T., Glacial geomorphology of the Gaumukh – the source of Ganga, with reference to its present state of environment. In Seminar on Ganga in the Service of the Nation, University of Roorkee, 12–13 September 1993, pp. 1–14.
17. Tangari, A. K., Chandra, R. and Yadav, S. K. S., Temporal monitoring of the snout, equilibrium line and ablation zone of Gangotri glacier through remote sensing and GIS techniques – an attempt at deciphering the climatic variability. In Proceedings of Workshop on Gangotri Glacier, Lucknow, 26–28 March 2003 (eds Srivastava, D., Gupta, K. R. and Mukerji, S.), Geological Survey of India Special Publication Number 80, 2004, pp. 145–153.
18. Bahuguna, I. M., Kulkarni, A. V., Nayak, S., Rathore, B. P., Negi, H. S. and Mather, P., Himalayan glacier retreat using IRS 1C PAN stereo data. *Int. J. Remote Sensing*, 2007, **28**, 437–442.
19. Kumar, R., Areendran, G. and Rao, P., Witnessing change: glaciers in the Indian Himalayas, WWF-Report (eds Mohan, D., Sinha, S. and Worah, S.), 2009, p. 48.
20. Bhambri, R. and Chaujar, R. K., Recession of Gangotri glacier (1962–2006) measured through remote sensing data. In Proceeding of National Seminar on Management Strategies for the Indian Himalaya: Development and Conservation, HNB Garhwal University, Srinagar, 2009, vol. 1, pp. 254–264.
21. Ahmad, S. and Hasnain, S. I., Analysis of satellite imageries for characterization of glacio-morphological features of the Gangotri Glacier, Ganga headwater, Garhwal Himalayas. In Proceedings of Workshop on Gangotri Glacier, Lucknow, 26–28 March 2003 (eds Srivastava, D., Gupta, K. R. and Mukerji, S.), Geological Survey of India Special Publication Number 80, 2004, pp. 61–67.
22. Vohra, C. P., Some problems of glacier inventory in the Himalayas. In Proceedings of the Workshop at Riederalp, 17–22 September 1978, IAHS-AISH Publ., 1980, vol. 126, pp. 67–74.
23. Agarwal, N. K., Remote sensing for glacier mapping and monitoring. Geological Survey of India Special Publication Number 53, 2001, pp. 201–206.
24. Raina, V. K. and Srivastava, D., *Glacier Atlas of India*, Geological Society of India, Bangalore, 2008, p. 316.
25. Raina, V. K., Himalayan glaciers: a state-of-art review of glacial studies, glacial retreat and climate change. Discussion Paper, Ministry of Environment and Forests, Government of India, 2009, p. 60.
26. Bhambri, R. and Bolch, T., Glacier mapping: a review with special reference to the Indian Himalayas. *Prog. Phys. Geogr.*, 2009, **33**(5), 672–704.
27. Paul, F. *et al.*, Recommendations for the compilation of glacier inventory data from digital sources. *Ann. Glaciol.*, 2009, **50**(53), 119–126.
28. Racoviteanu, A. E., Paul, F., Raup, B., Khalsa, S. J. S. and Armstrong, R., Challenges and recommendations in mapping of glacier parameters from space: results of the 2008 Global Land Ice Measurements from Space (GLIMS) workshop, Boulder, Colorado, USA. *Ann. Glaciol.*, 2009, **50**, 53–69.
29. Bolch, T., Menounos, B. and Wheate, R., Landsat-based glacier inventory of western Canada, 1985–2005. *Remote Sensing Environ.*, 2010, **114**(1), 127–137.
30. Bolch, T. *et al.*, A glacier inventory for the western Nyainqentanglha Range and Nam Co Basin, Tibet, and glacier changes 1976–2009. *Cryosphere*, 2010, **4**, 419–433.
31. Bhambri, R., Bolch, T. and Chaujar, R. K., Mapping of debris-covered glaciers in the Garhwal Himalayas using ASTER DEMs and thermal data. *Int. J. Remote Sensing*, 2011, **32**(23), 8095–8119.
32. Bolch, T., Buchroithner, M. F., Pieczonka, T. and Kunert, A., Planimetric and volumetric glacier changes in the Khumbu Himal, Nepal, since 1962 using Corona, Landsat TM and ASTER data. *J. Glaciol.*, 2008, **54**(187), 592–600.
33. Narama, C., Kääb, A., Duishonakunov, M. and Abdrakhmatov, K., Spatial variability of recent glacier area changes in the Tien Shan Mountains, Central Asia, using Corona (~1970), Landsat (~2000), and ALOS (~2007) satellite data. *Global Planet. Change*, 2010, **77**(1–2), 42–54.
34. Dashora, A., Lohani, B. and Malik, J. N., A repository of earth resource information – CORONA satellite programme. *Curr. Sci.*, 2007, **92**(7), 926–932.
35. Surazakov, A. and Aizen, V. B., Positional accuracy evaluation of declassified Hexagon KH-9 mapping camera imagery. *Photogramm. Eng. Remote Sensing*, 2010, **76**(5), 603–608.
36. Koblet, T., Gärtner-Roer, I., Zemp, M., Jansson, P., Thee, P., Haeberli, W. and Holmlund, P., Reanalysis of multi-temporal aerial images of Storglaciären, Sweden (1959–99) – Part 1: Determination of length, area, and volume changes. *Cryosphere*, 2010, **4**, 333–343.
37. Vohra, C. P., Gangotri Glacier. *Indian Mountaineer*, Indian Mountaineering Foundation, 1988, pp. 51–58.
38. Hall, D. K., Bahr, K. J., Shoener, W., Bindschadler, R. A. and Chien, J. Y. L., Consideration of the errors inherent in mapping historical glacier positions in Austria from the ground and space. *Remote Sensing Environ.*, 2003, **86**, 566–577.
39. Kargel, J. S., Cogley, J. G., Leonard, G. J., Haritashya, U., Byers, and Byers, A., Himalayan glaciers: the big picture is a montage. *Proc. Natl. Acad. Sci. USA*, 2011, **108**(36), 14709–14710.
40. http://asterweb.jpl.nasa.gov/content/03_data/05_application_examples/glacier/default.htm
41. Holmes, D. L., *Principles of Physical Geology*, Van Nostrand Reinhold (UK) Co Ltd, 1978, 3rd edn, p. 730.
42. Naithani, A. K., Nainwal, H. C., Sati, K. K. and Prasad, C., Geomorphological evidences of retreat of the Gangotri glacier and its characteristics. *Curr. Sci.*, 2001, **80**, 87–94.
43. Thayyen, R. J., Lower recession rate of Gangotri glacier during 1971–2004. *Curr. Sci.*, 2008, **95**, 9–10.
44. Bolch, T., Pieczonka, T. and Benn, D. I., Multi-decadal mass loss of glaciers in the Everest area (Nepal Himalaya) derived from stereo imagery. *Cryosphere*, 2011, **5**(2), 349–358.
45. Mukherjee, B. P. and Sangewar, C. V., Recession of Gangotri glacier through 20th century. Geological Survey of India Special Publication Number 65, 2001, pp. 1–3.

ACKNOWLEDGEMENTS. We thank the Director, Wadia Institute of Himalayan Geology (WIHG), Dehradun for support and Deepak Srivastava, CFG-WIHG, for thoughtful discussion on Gangotri Glacier monitoring programme by GSI and valuable suggestions. R.B. thanks Drs D. P. Dobhal and Manish Mehta, CFG-WIHG for support. We also thank Bhanu Pratap and Priyeshu Srivastava, CFG-WIHG for providing valuable suggestions and the anonymous reviewer for insightful comments and suggestions that helped improve the manuscript.

Received 25 April 2011; revised accepted 10 January 2012